

African Journal of Tropical Medicine and Biomedical Research (AJTMBR)



The Journal is the Official Publication of the College of Health Sciences, Delta State University, Abraka, Nigeria.

Editorial Board

Editor-in-Chief

Igbigbi P S

Nwagha E

Akpoyovwere OJ

Editor

Omo-Aghoja L O

Editorial Assistants

Adeyinka A T

Mofon CM

Associate Editors

Akhator A

Ejebe D E

Desk/Managing Editor

Aghedo O V

Editorial Advisory Board

Agbamu J U

Aloamaka C P

Asagba S O

Dosumu E A

Ebeigbe P N

Ekele B A

Fasubaa O B

Feyi-Waboso P

Ikomi R B

Obuekwe O N

Ohaju-Obodo J

Okobia M N

Okonofua F E

ISSN: 2141-6397

Focus and Scope

The *African Journal of Tropical Medicine and Biomedical Research* is a multidisciplinary and international journal published by the College of Health Sciences, Delta State University of Abraka, Nigeria. It provides a forum for Authors working in Africa to share their research findings on all aspects of Tropical Medicine and Biomedical Sciences and to disseminate innovative, relevant and useful information on tropical medicine and biomedical sciences throughout the continent. The journal will publish original research articles, reviews, editorials, commentaries, short reports, case reports and letters to the editor. Articles are welcome in all branches of medicine and dentistry including basic sciences (anatomy, biochemistry, physiology, pharmacology, psychology, nursing etc) and clinical sciences (internal medicine, surgery, obstetrics and gynaecology, dental surgery, child health, laboratory sciences, radiology, community medicine, etc). Articles are also welcome from social science researchers that document the intermediating and background social factors influencing health in countries of Africa. Priority will be given to publication of articles that describe the application of the principles of primary health care in the prevention and treatment of diseases.

Editorial Notices

The journal will be published biannually in the months of March and September. Annual subscription fee in Nigeria is two thousand naira (N2,000) per volume (2 issues); One thousand naira single copy (N1000). The annual subscription rate for other parts of the world is as follows: United Kingdom £60 (post free). West Africa \$60 (post free). The rest of the World and the United States of America \$120 (post free). A charge of \$60 is made for reprints inclusive of postage. Cheques should be made payable to the African Journal of Tropical Medicine and Biomedical Research and addressed to the Editor-in-Chief.

Journal Contact

All correspondence, including manuscripts for publication (in triplicate) should be addressed to:

Professor P.S. Igbigbi

The Editor-in-Chief,
Department of Anatomy,
Faculty of Basic Medical Sciences,
College of Health Sciences,
Delta State University,
Abraka, Delta State,
Nigeria.

Or:

Professor Lawrence Omo-Aghoja

Editor
Department of Obstetrics and
Gynecology,
Faculty of Clinical Medicine,
Delta State University,
Abraka, Nigeria.
Email: journal_ajtmbr@yahoo.com
Cc: all email to
eguono_2000@yahoo.com
Tel: 08039377043

All authors are advised to submit an electronic copy in CD-ROM along with a hard copy of their manuscript, as this will spare remarkable time in the reviewing and typesetting processes.

In the alternative, authors can submit their articles and covering letter by email attachments. A covering letter (signed by all authors) accompanying the manuscript should certify that the article has not been previously published and is not being considered for publication elsewhere.

Information for Authors

All manuscript are peer-reviewed and accepted with the understanding that the work has not been published or being considered for publication elsewhere. Indeed the authors would be requested to sign a copyright form transferring the ownership

of the paper to the *African Journal of Tropical Medicine and Biomedical Research*. All articles must include the correct names and addresses of author(s) including e-mail addresses and telephone numbers. Articles will be subjected to a thorough peer review process before any decision is made to publish or not. Authors should note that the *African Journal of Tropical Medicine and Biomedical Research* is not under any obligation to publish articles submitted, as decision to publish will be based on recommendations of reviewers and the editorial advisory board.

Manuscripts

Articles submitted for publication should be typed double-spaced with 2.5cm margins with accompanying CD-ROM in Microsoft Word format for easy and quick peer review and typesetting. Each of the following sections should begin in a new page: title page, abstract, introduction, materials and methods, results, discussion, acknowledgment (s), references, tables, legends to figures and illustrations. The manuscript should include:

Title Page

The title page should include the following information: 1. the title and sub-title; 2. the name(s) of the author(s); 3. the affiliation(s) of the author(s); 4. name and address of the corresponding author and 5. three to six key words for indexing and retrieval purposes.

Abstract

The abstract should be structured and not more than 250 words. It should carry the following headings: *Introduction, Materials and Methods, Results and Conclusion*.

Original Research - The journal welcomes articles reporting on original research, including both quantitative and qualitative studies. Full-length articles should generally not exceed 3000 words,

excluding abstract, tables, figures, and references. The subject matter should be organised under appropriate headings and sub-headings as itemized above.

Review Articles - Comprehensive review articles on all aspects of tropical medicine and biomedical sciences will also be considered for publication in the journal. Reviews should provide a thorough overview of the topic and should incorporate the most current research. The length of review articles must not exceed 3,000 words and the organisational headings and sub-headings used are at the author's discretion.

Short Reports - Brief descriptions of preliminary research findings or interesting case studies will be considered for publication as short reports. The length of the abstract and article should be restricted to 150 and 2,000 words respectively and organisation of short reports are left to the author's discretion.

Commentaries or Editorials - Commentaries or editorials on any aspect of tropical medicine and biomedical sciences in Africa will be considered for publication in the journal. Opinion pieces need not reference previous research, but rather reflect the opinions of the author(s). The length should not exceed 2,000 words.

Tables and Figures

All tables and figures should be submitted on separate sheets of paper and should be clearly labelled. Coloured tables and figures may be reprinted in black and white. Authors should especially take care that all tables are clear and understandable by themselves, independent of the text. A reader should be able to read only the tables and easily grasp all information without the text.

Acknowledgments

Acknowledgments should be included on a separate sheet of paper and should not exceed 100 words. Funding sources should be noted here.

References

References should be in the Vancouver style and numbered consecutively in the order in which they are mentioned in the text. Titles of journals should be abbreviated according to the *Index Medicus* style. Authors must cross-check and make sure that all information provided in the reference list is complete and correctly written. Reference numbers should be inserted above the line on each occasion a reference is cited in the text, e.g., ... as 1-3 reported in other studies. Numbered references should appear at the end of the article and should include the names and initials of all authors. The format of references should be as published by the International Committee of Medical Journal Editors in the British Medical Journal 1988, volume 296, pages 401-405. The following are sample references for an article published in a journal and for a book: Ahmed Y, Mwaba P, Chintu C, Grange JM, Ustianowski A, Zumla A. A study of maternal mortality at the University Teaching Hospital,

Lusaka, Zambia: the emergence of tuberculosis as a major non-obstetric cause of maternal death. *Int J Tuberc Lung Dis* 1999; 3: 675-680. Whitby LG, Smith AF, Beckett GJ. Enzyme Tests in Diagnosis. In: *Lecture Notes on Clinical Chemistry*. Whitby LG, Smith AF & Beckett GJth (eds). 4 edition. Blackwell Scientific Publications. 1988. 103-127.

Units of Measurement

All measurements should be expressed in SI (Systeme International) Units.

Galley proofs

Corrections of galley proofs should be strictly restricted to Printer's error only. Orders for offprints should be made when the corrected proofs are being returned by the authors. Articles accepted for publication remain the property of the journal and can only be reproduced elsewhere in line with section 5 of the copyright agreement.

Clinical Profile and Outcome of Adult Tetanus in Oghara, Delta State.

Obiabo Yo¹

ABSTRACT

Background: Tetanus persists to be a global problem due to either failed immunization or poor coverage of many inaccessible populations in both developing and developed societies. Mortality is higher in the developing world with poorly organized health systems.

Aim: To report the clinical profile and outcome of tetanus admissions in Delta State University Teaching Hospital, Oghara, between June 2010 and April 2015.

Methods: The records of all patients managed for tetanus on the medical wards and the intensive care unit of the hospital were studied.

Results: Twenty-seven patients comprising of 21(77.8%) males and 6 (22.2%) females were diagnosed and managed for tetanus during the period under review. The mean age of the patients was 34.07 ± 1.50 years. The patients were largely of low socioeconomic group. Lower extremity wounds and lesions was the commonest portal of entry, 74.1%. The overall case fatality rate (CFR) was 37%. Among those managed in the intensive care unit the case fatality was 50%. The factors significantly associated with increased mortality include short incubation period ($p = 0.009$), short onset time ($p = 0.001$), severity of the disease ($p = 0.014$) and increasing age ($p = 0.03$). The commonest complications recorded in this study were laryngeal and pharyngeal spasms, aspiration pneumonia and autonomic dysfunction.

Conclusion: Tetanus is life-threatening disease with high case fatality that is still a health challenge in our environment. There is need to provide better vaccination coverage for all including booster doses for adult males and to improve on medical intensive care facilities and training.

Keywords: *Clinical profile, outcome, Adult, Tetanus*

¹*Department of Internal Medicine, College of Health Sciences, Delta State University, Abraka, Nigeria*

Correspondence: *Obiabo YO, Department of Internal Medicine, College of Health Sciences, Delta State University, Abraka, Nigeria*

Introduction

Tetanus is a preventable life threatening neurological disorder that constitutes a major public health challenge in developing countries. It is a cause of significant morbidity and mortality especially in developing countries. Despite several concerted efforts through immunization to contain this disorder, it persists as a global problem due to either failed immunization or poor coverage of many inaccessible populations in both developing and developed societies.¹

Tetanus is caused by a neurotoxin produced by *Clostridium tetani*, a gram-positive, obligate anaerobic rod-shaped bacterium that forms spores. The spores are ubiquitous worldwide as

constituents of soil and in the gastrointestinal tracts of humans and animals whose waste can contaminate many surfaces and substances. The spores are extremely difficult to destroy requiring autoclaving or prolonged exposure to iodine, hydrogen peroxide, formalin or glutaraldehyde.² The exotoxin, tetanospasmin, which is one of the most potent toxins ever known to man, with a minimum lethal dose of less than 2.5 ng/kg in humans³ and absolute neurospecificity and enzymatic action,⁴ is responsible for the clinical manifestations. The Clinical manifestations are characterized by rigidity of muscles, painful muscle spasm, locked-jaw or trismus and clear mental

status.

Unhygienic practices such as use of old or unsterilized razor blade for making tattoos, tribal marks, cutting umbilical cords; close contact of broken skin or wounds with soil and animal dung,; unsafe abortions, lack of active and passive immunization are the major risk factors. This study described the clinical characteristics of adult tetanus in Delta State University Teaching hospital, Oghara, Nigeria.

Materials and Methods

We retrospectively reviewed the records of all patients with tetanus admitted into the hospital and managed by the neurology unit of Delta State University Teaching Hospital, Oghara from June 2010 to April 2015. Tetanus was clinically diagnosed based on the presence of at least two of the following features trismus, risus sardonicus, painful spasms and rigidity with or without opisthotonus. The demographic characteristics, portal of entry, incubation period, onset time, the severity, need for ventilator, duration of hospital stay, complications and outcome were obtained

and analyzed. Incubation period was defined as the time interval in days between injury sustained and the first symptom or sign of tetanus. The onset-time was defined as the time interval in days between the first symptom and the first spasm. The disease was categorized based on severity using the Ablett's scoring system⁵. The outcome of the admission was categorized into discharged or dead. Autonomic dysfunction was diagnosed if there were two or more of: labile blood pressure (hypertension and hypotension), high grade fever in the absence of infection, tachycardia/bradycardia and diaphoresis in the absence of fever. The data was analyzed using statistical package for social sciences (SPSS) version 21

Results

A total of 27cases were diagnosed and managed for tetanus out of 925 neurological admissions during the period under review giving a frequency of 2.92%. The patients comprised of 21 (77.80%) males and 6 (22.20%) females with a male to female ratio of 4.5: 1.0. See table 1.

Table 1: Age and Sex Distribution of Patients with Tetanus

Age group in years	< /=20	21-30	31-40	41-50	51-60	> 60	Freq. (%)
Males	3	6	8	0	2	2	21 (77.8)
Females	0	2	2	1	0	1	6(22.2%)
Total	3	8	10	1	2	3	27 (100%)

The mean age of the patients was 34.07 ± 1.50 years with age range of 13years to 65 years. The mean age for males was 33.04 ± 1.50 years and mean age for females was 37.67 ± 1.52 years. The patients were made up of artisans 9(33.3%), traders and business men and women 6(22.2%), farmers 4(14.8%), students 4(14.8%), civil servants 3(11.1%) and retired personnel 1(3.7%). The portal of entry was puncture wounds and lacerations (including 2 cases of machet cuts) in 24 (88.9%), diabetic foot ulcer was recorded in a patient(3.7%) and no

obvious portal of entry was found in 2 patients (7.4%). Lower extremity lesions constitute portal of entry in 20 (74.1%) of cases.

The mean incubation period was 6.70 ± 2.83 days with a range of 3 to 14 days. The mean onset time was 2.04 ± 0.85 days, ranging from 1 day to 4 days. Those who survived had longer incubation period of 7.6 ± 3.1 days compared to those who died with incubation period of 5.2 ± 1.3 days. The onset time was also shorter

among those who died (1.2 ± 0.42 days) compared to those who survived with onset time of 2.5 ± 0.62 days. The mean dose of diazepam received was $216.2\text{mg} \pm 112.04\text{mg}$ per day. There was no significant difference between the mean dose received by those managed in the intensive care unit and those managed in the open ward $p=0.234$.

All the patients (100%) presented with lock-jaw and painful spasms while 25 (92.6%) had risus sardonicus. Rigidity with opisthotonus posturing was recorded in 22 (81.50%) and 20 (74.10%) had board-like rigidity of the abdomen. See table 2.

The severity grading showed that none (0%) had mild tetanus, 4 (14.8 %) had moderate, 9 (33.3 %) had severe and 14 (51.9%) had very severe tetanus. Generalised tetanus was present in 26 (96.3%) and localized tetanus was seen in only one patient (3.7%). See table 2.

Out of the 27 patients 17 (63%) survived and were discharged home while 10 died constituting case fatality rate of 37%. The mean age for the survivors was 31.3 ± 1.3 years while the mean age for those who died was 38.6 ± 1.7 years. For those that were managed in the intensive unit, the case fatality rate

Table 2: Presenting Clinical Features & Complications of Tetanus among the Patients

Clinical features	Frequency	Percentage (%)
Portal of Entry		
Puncture wounds & lacerations	24	88.9
Diabetic foot ulcer	1	3.7
No obvious portal	2	7.4
Presenting symptoms & signs		
Trismus	27	100
Painful spasms	27	100
Neck rigidity & Opisthotonus	22	81.5
Risus Sardonicus	25	92.6
Abdominal rigidity (board like)	20	74.1
Complications		
Laryngeal/Pharyngeal Spasms	14	51.9
Aspiration pneumonia	11	40.7
Sepsis	5	18.5
Acute kidney injury	2	7.4
Autonomic dysfunction	8	29.6
DVT	1	3.7
Ablett's Severity grade		
I Mild	0	0
II Moderate	4	14.8
III Severe	9	33.3
IV Very Severe	14	51.9

DVT = Deep vein thrombosis

was 50%. Out of the twenty-one patients who were managed in the emergency room or the wards there were 7 deaths with a case fatality rate of 33.3%. The cause of death of those who died in the emergency room or on the ward was mainly from respiratory failure secondary to laryngeal spasm.

The factors significantly associated with increased mortality included short incubation period less than 7 days ($p = 0.009$), short onset time less than 2 days ($p = 0.001$), severity of the disease ($p = 0.014$) and increasing age ($p = 0.03$). See table 3. Those with bad outcome tended to have shorter duration of hospital stay but there was no significant

difference between severity of the disease and the duration of hospital stay ($p = 0.37$). Also there was no significant difference between the sexes with respect to outcome.

Discussion

This study showed that despite concerted efforts to curb the burden of the disease through immunization coverage in the country, tetanus is still a cause of morbidity and mortality in Nigeria. Tetanus tend to occur in the younger working population in developing countries compared to st

Table 3. Comparison of mean variables between outcomes

Variable	Discharges	Deaths	p-value
Age (years)	31.30±1.30	38.6±1.70	0.030
Incubation period (days)	7.60±3.10	5.20 ±1.30	0.009
Onset time	2.50±0.62	1.20±0.42	0.001
Duration of hospital stay	23.76±1.20	7.10±6.24	0.370

the developed nations.¹⁷ The mean age of occurrence in our study was 34.1 ± 1.5 years. This supports the observations of most studies in the country and outside the country.⁶⁻¹⁰ Komolafe et al in Ife however, found a mean age of 53 years in their patient population.¹¹ The frequency of tetanus in this survey is lower than most studies in the country.⁹⁻¹³ This is likely a reflection of immunization coverage. A lower frequency of tetanus was recorded among females with a male to female ratio of 4.5: 1. This has also been reported in some previous local and foreign studies.⁶⁻¹⁷ There is no known biological reason in males or in the pathobiology of the disease that can explain the sex difference. The observed discrepancy may be linked to lack of protective immunity against tetanus occasioned by lack of booster doses of the tetanus vaccination among men in comparison to the improved immunization coverage with booster doses for women especially during pregnancy.^{17,18} Ogunrin et al however, reported slight

preponderance in females.¹⁹ The reason for this finding is not well understood.

The patients in this survey were largely from the lower socioeconomic strata of the society and the portal of entry for the disease was largely lower extremity injuries and lesions. The commonest clinical manifestation of tetanus among them were trismus and painful spasms (100%) followed by rigidity of the neck and risus sardonicus. This clinical profile has similarly been reported in the literature by several authors.⁶⁻¹⁶

Tetanus is a dreaded disease with records of high mortality. The overall case fatality rate (CFR) from this study was 37%. Ojini and Danesi reported a similar case fatality rate of 36.96% in a study of 349 cases between 1990 and 1999 from Lagos,¹⁵ but Sanya et al reported 64% in a review of 202 cases between 1990 and 2001 from Ibadan²⁰ and in Benin City, Ogunrin et al reported a mortality rate of 26.2% with an age-adjusted fatality rate of 16.2% for those less than 40 years of age in a review of 66

cases between 1990 and 2000.¹⁹ Higher mortality rates of 57.1%, 53.5%, 44% were also reported from in southwest Nigeria by different authors.^{7,11,16.}

Owolabi et al reported mortality of 46% from Kano in a 10-year review⁸ whereas Chukwubike and God'spower reported a case fatality rate of 42.9% among 86 cases from PortHarcourt.⁶ In Ethiopia Amanuel et al reported a case fatality rate of 35.3%¹³ whereas in India, Anuradha reported 37.78%.²¹

Intensive care facilities have been found to reduce mortality from tetanus and mortality rates as low as 10% have been reported from units with facilities for intensive care.^{23,24} The case fatality recorded in this study is however 50% percent mortality among the patients managed in the intensive care unit. One of the mortality cases in the intensive care unit was caused by failure to regain consciousness from the effect of anaesthesia after being weaned off from ventilation two weeks after cessation of spasms. The second patient died from overwhelming sepsis and ensuing renal failure. The third patient developed severe hypotension following intubation and died of cardiac arrest. Interestingly one patient managed in the intensive care unit had tracheostomy without ventilator support and performed excellently well despite the presence of complications. It is not very clear why intensive care services did not improve the outcome as expected among the patients. Plausible reasons include severe hypotension induced by induction agents and oversedation. Ojini et al also found that intensive care did not reduce mortality.¹⁵

The poor prognostic indicators in our study included short incubation period less than 7 days, onset time less than 2 days, severe tetanus especially with autonomic features and the presence of complications like aspiration pneumonia, severe sepsis and acute kidney. These observations have been documented in other studies.^{15,16,19,21}

No case of post-abortal or puerperal tetanus was recorded during the period under review. This is in sharp contrast to findings in previous studies¹¹ and may also reflect an improvement in maternal and child health care services in recent years in the region.

The complications commonly seen among our patients included aspiration pneumonia, laryngeal spasm and respiratory failure, autonomic dysfunction with marked diaphoresis, labile hypertension, arhythmias (tachy- or bradycardia), acute kidney injury and sepsis. These observations support the findings in both local and foreign studies.^{8-16,19}

The specific objectives of tetanus treatment are to stop the production of toxin at the site of infection with appropriate wound care and antibiotic use; to neutralize circulating toxin with anti-tetanus immunoglobulin; and to provide effective management of muscle spasm, respiratory failure, autonomic dysfunction, and complications. The patients in our study received wound care where present and metronidazole with or without any other antibiotic, anti-tetanus serum /immunoglobulin and high doses of diazepam to control spasms. The average dose of diazepam received was 216.2mg ± 112.04mg per day. The use of these treatment modality have been reported to reduce mortality and morbidity of tetanus patients admitted to hospital.¹⁵ Even in settings with limited resources, if basic medication, experienced medical supervision, and high-quality nursing can be provided, mortality can be reduced.²⁵ In the author's experience the greatest impediment to improved survival of tetanus patients in our environment is the lack of access to appropriate medical care.

Conclusion

Tetanus is a preventable life threatening illness that is still a significant cause of morbidity and mortality in Delta State as observed in this hospital, especially among young working class males in their prime of life. There is need to increase efforts at awareness creation and providing protective tetanus immunization especially for men in form of booster doses during preschool and pre-employment medical examination programs. Also improvement in hygienic practices and increased nursing care and medical supervision will go a long way to reduce the attendant mortality in patients with tetanus.

References

1. Centers for Disease Control and Prevention. Tetanus. In W Atkinson et al., eds. *Epidemiology and Prevention of Vaccine-Preventable Diseases*, 12th ed., Washington, DC: Public Health Foundation. 2011, pp. 291300.
2. Feingold SM. Tetanus. In: Collier L, Balows A, Sussman M, eds. *Topley & Wilson's Microbiology and Microbial Infections*. 9th ed. New York: Oxford University Press, 1998: 694722.
3. Gill DM. Bacterial toxins: a table of lethal amounts. *Microbiol Rev* 1982; 46: 8694.
4. Schiavo G, Matteoli M, Montecucco C. Neurotoxins affecting neuroexocytosis. *Physiol Rev* 2000; 80: 717766.
5. Ablett JJJ. Analysis and main experiences in 82 patients treated in the Leeds Tetanus Unit. In: Ellis M, editor. *Symposium on tetanus in Great Britain*. Boston Spa, UK: Leeds General Infirmary; 1967. p. 1-10.
6. Chukwubike OA, God'spower AE. A 10-year review of outcome of management of tetanus in adults at a Nigerian tertiary hospital. *Ann Afr Med* 2009; 8: 168-172.
7. Fawibe AE. The Pattern and Outcome of Adult Tetanus at a Sub-urban Tertiary Hospital in Nigeria *Journal of the College of Physicians and Surgeons Pakistan* 2010, Vol. 20 (1): 68-70
8. Owolabi LF, Aliyu I, Nagoda, M et al. Clinical profile and outcome of adult tetanus in Kano. *Sahel Medical Journal*, 2010, Vol 13, (4):67-68. <http://dx.doi.org/10.4314-smj2.v13i4.67508>
9. Tadessa A, Gebre-Selassie S. Five year review of cases of adult tetanus management at Gondar University Hospital, North West Ethiopia (Gondar, Sept.2003-Aug 2008). *Ethiop Med. J.* Oct 2009; 47(4): 291-7.
10. Onwuekwe IO, Nwabueze AC. Experience with tetanus in a tertiary hospital in the South East Nigeria. *Nigerian Journal of Medicine journal of Resident doctors of Nigeria*. April 2008; 17(1): 50-52
11. Komolafe MA, Komolafe EO, Oqundere AO. Pattern and outcome of adult tetanus in Ile-Ife, Nigeria, *Niger J. Clinical. Practice*. Dec 2007; 10 (4): 300-3.
12. Adeuja AOG, Osuntokun BO. Tetanus in the adult Nigerians. A review of 503 patients . *East . Africa Med.*, 1971; 4(12): 683-691
13. Amanuel A, Yilma M, Desalar M. Tetanus in adults : clinical presentation, treatment and mortality in Ethiopia. *Journal of Neurological sciences*. Mar 2012; 317(1-2): 62-65
14. Sinha A, Seth BC, Ali MH, Mondal K, Bhattacharya R .Study of Clinical Profile of Tetanus Patients at ID & BG Hospital, Kolkata. *National Journal of Medical and Allied Sciences* 2014; 3(2):4-8
15. Ojini FI, Danesi MA Mortality of tetanus at the Lagos University Teaching Hospital, Nigeria. *Top Doct*, 2005 Jul; 35(3):178-11
16. Arogudade FA Bello IS, Kuteyi EA, Akinsola A. Patterns of presentation and mortality in tetanus: a 10-year retrospective review. *Niger Postgrad Med J*. 2004 Mar 11(1); 58-63
17. Centers for Disease Control and Prevention. Advisory Committee on Immunization Practices (ACIP) Recommended Immunization Schedules for Persons Aged 0 Through 18 Years and Adults Aged 19 Years and Older. United States, 2013. *MMWR*. 2013; 62(Suppl 1):1-19.
18. Orimadegwu AE, Adepoju AA, Akinyinka OO .Prevalence and social demographic factors associated with non-protective immunity against tetanus among high school adolescent girls in Nigeria. *Ital J. Paediatr* 2014 , Mar 17; 40(1): 29-32 . do; .10.118/1824-7288-40-29
19. Ogunrin OA, Unuigbo EI Tetanus: an analysis of prognosticating factors of cases admitted into the medical wards of a tertiary hospital in a developing African country between 1990 and 2000. *Niger Postgrad Med J*, 2004 Jun; 11(2):9-102
20. Sanya EOI, Taiwo SS, Olarinoye et al. A 12-year review of cases of adult tetanus

- managed at the University College Hospital, Ibadan, Nigeria. Trop Doct. 2007 Jul; 37(3): 170-173.
21. Anuradha S. Tetanus in adults, a continuing problem: Analysis of 217 patients over 3 years from Delhi, India, with special emphasis on predictors of mortality. Med J Malaysia. 2006 Mar;61(1):7-14.
 22. Bardenheier B, Prevots DR, Khetsuriani N et al. Tetanus surveillance- united States, 1995-1997: Morbidity and Mortality Weekly Report; 1998. Report No. 47: p. 1-13.
 23. Brauner JS, Vieira SR, Bleck TP. Changes in severe accidental tetanus mortality in the ICU during two decades in Brazil. Intensive Care Med 2002; 28: 93035.
 24. Thwaites CL, Yen LM, Nga NT, et al. Impact of improved vaccination programme and intensive care facilities on incidence and outcome of tetanus in southern Vietnam, 1993-2002. Trans R Soc Trop Med Hyg 2004; 98: 67177.
 25. Salimpour R. Tetanus of the newborn in Tehran. A ten year study of 880 cases. J Trop Pediatr Environ Child Health 1978; 24: 14042.

Citation

This article should be cited as: "Obiabo YO. *Clinical Profile and Outcome of Adult Tetanus in Oghara, Delta State*. Afr. J. Trop. Med. & Biomed. Res 2014; 3 (1): 21-27".