

# African Journal of Tropical Medicine and Biomedical Research (AJTMBR)



The Journal is the Official Publication of the College of Health Sciences, Delta State University, Abraka, Nigeria.

## *Editorial Board*

### **Editor-in-Chief**

*Igbigbi P S*

*Nwagha E*

*Akpoyovwere OJ*

### **Editor**

*Omo-Aghoja L O*

### **Editorial Assistants**

*Adeyinka A T*

*Mofon CM*

### **Associate Editors**

*Akhator A*

*Ejebe D E*

### **Desk/Managing Editor**

*Aghedo O V*

## **Editorial Advisory Board**

*Agbamu J U*

*Aloamaka C P*

*Asagba S O*

*Dosumu E A*

*Ebeigbe P N*

*Ekele B A*

*Fasubaa O B*

*Feyi-Waboso P*

*Ikomi R B*

*Obuekwe O N*

*Ohaju-Obodo J*

*Okobia M N*

*Okonofua F E*

ISSN: 2141-6397

### **Focus and Scope**

The *African Journal of Tropical Medicine and Biomedical Research* is a multidisciplinary and international journal published by the College of Health Sciences, Delta State University of Abraka, Nigeria. It provides a forum for Authors working in Africa to share their research findings on all aspects of Tropical Medicine and Biomedical Sciences and to disseminate innovative, relevant and useful information on tropical medicine and biomedical sciences throughout the continent. The journal will publish original research articles, reviews, editorials, commentaries, short reports, case reports and letters to the editor. Articles are welcome in all branches of medicine and dentistry including basic sciences (anatomy, biochemistry, physiology, pharmacology, psychology, nursing etc) and clinical sciences (internal medicine, surgery, obstetrics and gynaecology, dental surgery, child health, laboratory sciences, radiology, community medicine, etc). Articles are also welcome from social science researchers that document the intermediating and background social factors influencing health in countries of Africa. Priority will be given to publication of articles that describe the application of the principles of primary health care in the prevention and treatment of diseases.

### **Editorial Notices**

The journal will be published biannually in the months of March and September. Annual subscription fee in Nigeria is two thousand naira (N2,000) per volume (2 issues); One thousand naira single copy (N1000). The annual subscription rate for other parts of the world is as follows: United Kingdom £60 (post free). West Africa \$60 (post free). The rest of the World and the United States of America \$120 (post free). A charge of \$60 is made for reprints inclusive of postage. Cheques should be made payable to the African Journal of Tropical Medicine and Biomedical Research and addressed to the Editor-in-Chief.

### **Journal Contact**

All correspondence, including manuscripts for publication (in triplicate) should be addressed to:

**Professor P.S. Igbigbi**

The Editor-in-Chief,  
Department of Anatomy,  
Faculty of Basic Medical Sciences,  
College of Health Sciences,  
Delta State University,  
Abraka, Delta State,  
Nigeria.

Or:

**Professor Lawrence Omo-Aghoja**

Editor  
Department of Obstetrics and  
Gynecology,  
Faculty of Clinical Medicine,  
Delta State University,  
Abraka, Nigeria.  
Email: journal\_ajtmbr@yahoo.com  
Cc: all email to  
eguono\_2000@yahoo.com  
Tel: 08039377043

All authors are advised to submit an electronic copy in CD-ROM along with a hard copy of their manuscript, as this will spare remarkable time in the reviewing and typesetting processes.

In the alternative, authors can submit their articles and covering letter by email attachments. A covering letter (signed by all authors) accompanying the manuscript should certify that the article has not been previously published and is not being considered for publication elsewhere.

### **Information for Authors**

All manuscript are peer-reviewed and accepted with the understanding that the work has not been published or being considered for publication elsewhere. Indeed the authors would be requested to sign a copyright form transferring the ownership

of the paper to the *African Journal of Tropical Medicine and Biomedical Research*. All articles must include the correct names and addresses of author(s) including e-mail addresses and telephone numbers. Articles will be subjected to a thorough peer review process before any decision is made to publish or not. Authors should note that the *African Journal of Tropical Medicine and Biomedical Research* is not under any obligation to publish articles submitted, as decision to publish will be based on recommendations of reviewers and the editorial advisory board.

### **Manuscripts**

Articles submitted for publication should be typed double-spaced with 2.5cm margins with accompanying CD-ROM in Microsoft Word format for easy and quick peer review and typesetting. Each of the following sections should begin in a new page: title page, abstract, introduction, materials and methods, results, discussion, acknowledgment (s), references, tables, legends to figures and illustrations. The manuscript should include:

### **Title Page**

The title page should include the following information: 1. the title and sub-title; 2. the name(s) of the author(s); 3. the affiliation(s) of the author(s); 4. name and address of the corresponding author and 5. three to six key words for indexing and retrieval purposes.

### **Abstract**

The abstract should be structured and not more than 250 words. It should carry the following headings: *Introduction, Materials and Methods, Results and Conclusion*.

Original Research - The journal welcomes articles reporting on original research, including both quantitative and qualitative studies. Full-length articles should generally not exceed 3000 words,

excluding abstract, tables, figures, and references. The subject matter should be organised under appropriate headings and sub-headings as itemized above.

Review Articles - Comprehensive review articles on all aspects of tropical medicine and biomedical sciences will also be considered for publication in the journal. Reviews should provide a thorough overview of the topic and should incorporate the most current research. The length of review articles must not exceed 3,000 words and the organisational headings and sub-headings used are at the author's discretion.

Short Reports - Brief descriptions of preliminary research findings or interesting case studies will be considered for publication as short reports. The length of the abstract and article should be restricted to 150 and 2,000 words respectively and organisation of short reports are left to the author's discretion.

Commentaries or Editorials - Commentaries or editorials on any aspect of tropical medicine and biomedical sciences in Africa will be considered for publication in the journal. Opinion pieces need not reference previous research, but rather reflect the opinions of the author(s). The length should not exceed 2,000 words.

### **Tables and Figures**

All tables and figures should be submitted on separate sheets of paper and should be clearly labelled. Coloured tables and figures may be reprinted in black and white. Authors should especially take care that all tables are clear and understandable by themselves, independent of the text. A reader should be able to read only the tables and easily grasp all information without the text.

### **Acknowledgments**

Acknowledgments should be included on a separate sheet of paper and should not exceed 100 words. Funding sources should be noted here.

## References

References should be in the Vancouver style and numbered consecutively in the order in which they are mentioned in the text. Titles of journals should be abbreviated according to the *Index Medicus* style. Authors must cross-check and make sure that all information provided in the reference list is complete and correctly written. Reference numbers should be inserted above the line on each occasion a reference is cited in the text, e.g., ... as 1-3 reported in other studies. Numbered references should appear at the end of the article and should include the names and initials of all authors. The format of references should be as published by the International Committee of Medical Journal Editors in the British Medical Journal 1988, volume 296, pages 401-405. The following are sample references for an article published in a journal and for a book: Ahmed Y, Mwaba P, Chintu C, Grange JM, Ustianowski A, Zumla A. A study of maternal mortality at the University Teaching Hospital,

Lusaka, Zambia: the emergence of tuberculosis as a major non-obstetric cause of maternal death. *Int J Tuberc Lung Dis* 1999; 3: 675-680. Whitby LG, Smith AF, Beckett GJ. Enzyme Tests in Diagnosis. In: *Lecture Notes on Clinical Chemistry*. Whitby LG, Smith AF & Beckett GJth (eds). 4 edition. Blackwell Scientific Publications. 1988. 103-127.

## Units of Measurement

All measurements should be expressed in SI (Systeme International) Units.

## Galley proofs

Corrections of galley proofs should be strictly restricted to Printer's error only. Orders for offprints should be made when the corrected proofs are being returned by the authors. Articles accepted for publication remain the property of the journal and can only be reproduced elsewhere in line with section 5 of the copyright agreement.

# Table of Contents

## Editorial

- Male Factor Infertility in Low Resource Settings: Diagnostic and Treatment Barriers 6  
*Omo-Agboja LO*

## Original Articles

- Acute Kidney Injury in Adult Nigerians: A Single Centre Experience 9  
*Okoye O, Unuigbo E, Ojogwu L.*
- Limb amputations at a Tertiary Hospital in the Developing World:  
A Look at Pattern and Indications 16  
*Bafor A , Akinyemi O*
- Clinical Profile and Outcome of Adult Tetanus in Oghara, Delta State. 21  
*Obiabo YO*
- Prevalence of Cardiovascular Risk Factors Among Factory Workers in the  
Beverage Industries in Edo State, Nigeria 28  
*Awunor NS, Isah EC*
- Patterns of Hysterosalpingographic Findings at a Tertiary Hospital in South-South,  
Nigeria: A Three-Year Review 34  
*Joyce E. Ikubor and Godson U. Eze*
- Assessing Reliability of Card Agglutination Test for Trypanosomiasis in Screening  
Gambian Sleeping Sickness at Abraka Endemic Focus 41  
*Umukoro DO, Ataikiru PD*
- Sputum AFB Positivity of Suspected Tuberculosis Patients in a Tertiary  
Hospital in Benin City Nigeria 46  
*Osarogiabon WO, Iyoha O*

# Editorial

## Male Factor Infertility in Low Resource Settings: Diagnostic and Treatment Barriers

Omo-Aghoja LO

Department of Obstetrics and Gynecology, Faculty of Clinical Medicine, College of Health Sciences, Delta State University, Abraka, Nigeria

**Keywords:** Male factor infertility, low resource settings, Diagnostic, Treatment, Barriers

**Correspondence:** Department of Obstetrics and Gynecology, Faculty of Clinical Medicine, College of Health Sciences, Delta State University, Abraka, Nigeria. Email: [eguono\\_2000@yahoo.com](mailto:eguono_2000@yahoo.com) Tel: +2348039377043

### Background

Male infertility refers to the inability of a male to achieve pregnancy in a fertile female. Hitherto in most part of the world particularly in Africa and other developing countries, women, bore the sole blame for barren marriages and took the psychosocial brunt; and in many areas infertility is a socially acceptable basis for divorce by the husband.<sup>1,3</sup> This may be due to the believe and inadequate knowledge and clear understanding existing in the sociocultural milieu of the couples. Male infertility is considered when identifiable female causes of infertility are excluded and semen quantity and quality fails to fulfill WHO criteria<sup>4,5</sup>

From available indices, Male factor infertility is quite common in developing nations and low resource settings accounting for 40-50% of infertility in humans,<sup>6-8</sup> and the most severe form of male infertility is that due to azoospermia which is present in 5% of investigated infertile couples<sup>9</sup>, this is against the backdrop of the fact that socio-culturally this is down played in most of these settings even in face of clear evidence that the male may be sole cause of the infertility.<sup>1,2,3</sup> Indeed in some settings the concealment of the male factor may lead to family members coercing the women to having sexual intercourse with a brother or other male relations of the husband to get her pregnant on behalf of the husband. This sociocultural perspectives no doubt significantly impacts on the incidence of male infertility which may indeed be higher than the currently quoted figures and the

treatment outcomes for those established cases, as compliance and follow-up of treatment is usually poor on the part of the males. The fact that males of infertile couples may not present for evaluation and seminal fluid analysis, continue to give relevance to the SIMs Huhner's test in developing countries/low resource settings.<sup>10,11</sup>

Beyond the socio-cultural impediments that continue to remain a major challenge to the management of male infertility in developing countries and low resource settings, availability and affordability of investigations for male infertility, treatment facilities and expertise for management remains poor.

### Discussion

There is paucity of health care facilities with the requisite needs for handling male infertility. Largely, the primary and secondary healthcare facilities that are closer to the greater majority of the inhabitants of low resource settings in these nations are devoid of these management needs. The tertiary health care facilities with laboratories where some of the modalities for investigation such as the seminal fluid analysis and the presence of a Gynecologist may not be accessible and affordable by a larger proportion of couples

with male factor infertility. They may be discouraged by the distance from their place of residence, lack of affordable means of transportation and the often prohibitive cost of treatment.

Hormone assays, facilities for genetic studies including karyotyping, testicular ultrasonography, vasogram and testicular biopsy are usually not readily available even in tertiary centers in most developing countries and low resource settings. Where they are available, they may not be readily accessed or afforded by the clients suffering male infertility for reason of cost as most patients with this condition are usually poor and there is no support for management of patients with infertility in countries of the globe.<sup>23,13</sup>

Again the needed experts for the full complements of investigations and treatment modalities such as a geneticist and urologists are usually not readily available even in tertiary health care facilities in developing countries. This further accentuates the difficulty to access the only few distant centers from the abode of clients with male factor infertility and the cost becomes usually more prohibitive because of paucity in face of high demand.

Poor health seeking behavior of citizens of most developing nations also challenges male factor infertility. Some infertile couples often consult spiritual homes, herbalists, traditional healers before going to hospital. This is because of perceived spiritual causes of infertility. When they finally present to orthodox practitioners, they tend to move from one practitioner to another. Sometimes they go back to traditional healers because of the high cost of investigations and treatment for infertility.<sup>10,11</sup>

Continuous search for treatment of male infertility particularly the variants with etiological consideration being testicular factors (which are not readily amenable to conventional therapy), led to the discovery of in-vitro fertilization (IVF) and other assisted conception techniques. This option is very expensive and not readily accessible in

developing countries. Extremely few public health care facilities have units for IVF and assisted conception, rather it is mainly private sector driven. The cost even in the few public health care facilities is usually prohibitive and out of the reach of very many clients. The acceptability of IVF from lack of appropriate information, impediments based on cultural and religious perspectives is another major challenge faced by this treatment option and has in some instances limited its use even by some that could have afforded it.

Adoption still remains a viable option for some couples where all other options fail. However this is poorly utilized in most developing countries. There are yet no clear laws and guidelines for its practice in most developing nations. The practice is further challenged in low resource settings and developing countries by serious sociocultural resentment to it.<sup>14</sup>

Poor budgetary allocations and or non-implementation of the meager allocation to health in face of poor political commitments to health care needs are other serious challenges in low resource settings. This makes the health care facilities poorly equipped in terms of facilities and requisite personnel.<sup>12</sup> The case for infertility management is made worse because of the strong moral and ethical debate against funding of infertility management with public funds in most countries of the world including developing countries.<sup>12</sup>

Finally, a large proportion of the citizenry in developing nations live in resource poor settings where there are no access to health care facilities, and for those that even live close to the health care facilities, they may not have the economic wherewithal to afford hospital care. Therefore poor personal and individual state of finance remains yet another major impediment and challenge to accessing health care facilities to seek treatment for male infertility. The policy of health care delivery in most developing countries is based on a "cash and carry" principle and in the face of ravaging poverty this will serve as a significant impediment.

### Conclusion

In conclusion, male factor contributes significantly to infertility globally with developing nations worse hit with its effect. The diagnosis and treatment of male infertility in low resources settings is significantly bedeviled with a number of intermediating socio-cultural, religious, economic and political challenges. Using conventional methods and recent technologies are expensive and not readily available in these environments. It is therefore necessary for us to understand the various factors that contribute to male infertility in our environment so as to develop preventive strategies.

### References

1. Singh, A.J. Support for infertile couples. *World Health Forum* 17:176-177(1996)
2. Leke, R.J.I. et al. Regional and geographic variations in infertility: Effects of environmental, cultural, and socioeconomic factors. *Environmental Health Perspectives Supplement* 101(Suppl. 2):73-80 (1993).
3. Yeboah, E.D. et al. Etiological factors of male infertility in Africa. *International Journal of Fertility* 37(5):300-307 (1992);
4. World Health Organization. WHO Laboratory manual for the examination and processing of human semen, 5 edition. Cambridge: Cambridge University Press, 2010;
5. Olooto, W.E. Infertility in male; risk factors, causes and management- A review *J. Microbiol. Biotech. Res.*, 2012, 2 (4):641-645.
6. "Men's Health - Male Factor Infertility". University of Utah Health Sciences Center. 2003-04-01. Archived from the original on 2007-07-04. Retrieved 2007-11-21;
7. Brugh VM, Lipshultz LI (2004). "Male factor infertility". *Medical Clinics of North America* 88(2):36785. doi :10.1016/S0025-7125(03)00150-0; A
8. Hirsh. Male subfertility, *BMJ*, 2003, 327 (7416): 669672.
9. Irvine DS. Epidemiology and etiology of male infertility. *Hum Reprod* 1998; 13 (soppl 1): 33-44.
10. Nwajiaku LA ,Mbachu II , Ikeako L. Prevalence, Clinical Pattern and Major Causes of Male Infertility in Nnewi, South East Nigeria: A Five Year Review. *AFRIMEDIC Journal* 2012; 3: 16-19;
11. Idrisa A. Infertility. In: Kwawukume EY, Emuveyan EE (Eds). *Comprehensive Gynaecology in the tropics*. Accra Graphic packaging, 2005:333-343
12. Oluyemi Akinloye and Ernest J Truter . A review of management of infertility in Nigeria: framing the ethics of a national health policy. *Int J Womens Health*. 2011; 3: 265275.
13. Philip Teg-Nefaah Tabong<sup>1</sup> and Philip Baba Adongo<sup>2</sup>. Infertility and childlessness: a qualitative study of the experiences of Infertile couples in Northern Ghana. *BMC Pregnancy and Childbirth* 2013, 13:72.
14. Oladokun A, Arulogun O, Oladokun R, Moharson-Bello IO, Bamgboye EA, Adewole IF et al. Acceptability of child Adoption as management option for infertility in Nigeria: Evidence from Focus Group Discussions. *Afr. J. Reprod. Health* 2009; 13 [1]: 79-91.